

CYGNUS (CYprus models for Galaxies and their NUClear Spectra)

Polar dust models

Andreas Efstathiou

Version 1.1

March 2018

The polar dust models have been used for the interpretation of the spectral energy distribution of the hyperluminous infrared galaxy IRAS F10214+4724 in Efstathiou (2006) and Efstathiou et al (2013). They have also been used for the interpretation of the tidal disruption event in Arp299 by Mattila et al (2018). The models form part of the CYGNUS collection of models (<http://ahpc.euc.ac.cy/index.php/resources/cygnus>).

A small library of models is included in the IDL saveset 'polar_dust_library1.dat'. The saveset includes the structure POLAR_DUST with the following fields:

```
TEMPERATURE  FLOAT    400.000  
WAVELENGTH   FLOAT    Array[223]  
NUSNU        FLOAT    Array[223]
```

The temperature is in Kelvin, the wavelength is in microns and nuSnu is normalized to give a bolometric luminosity of unity. The library of models can be used to synthesize an ensemble of clouds in the polar region with a range of temperature and covering factors.

Enquiries should be addressed to Andreas Efstathiou (a.efstathiou@euc.ac.cy).

References

Efstathiou, A., 2006, MNRAS, 371, L70
Efstathiou, et al., 2013, MNRAS, 436, 1873
Mattila et al., 2018, to appear in Science